Canberra heart clinic	
Right Heart Catheter (Interna Jugular Vein Approach)	al
A. Interpreter / cultural needs	
An Interpreter Service is required?	s [
If Yes, is a qualified Interpreter present?	s [
A Cultural Support Person is required?	s [
If Yes, is a Cultural Support Person present?	- [

B. Condition and treatment

The doctor has explained that you have the following condition: (Doctor to document in patient's own words)

This condition requires the following procedure. (Doctor to document - include site and/or side where relevant to the procedure)

The following will be performed:

The vein in your neck is found with ultrasound.

You will be given an injection of local anaesthetic. A fine tube (catheter) is put into the vein in the neck. You may feel pressure on your neck while the tube is placed in the vein. It is passed down until it reaches the heart and then goes up into the blood vessels of the lungs. This is usually painless. The doctor uses x-ray imaging to see the catheter.

Pressures in the lungs and the heart are recorded. A sample of blood is taken to look at the oxygen level and other levels.

At the end of the procedure, the catheter and sheath are removed.

C. Risks of a right heart catheter (internal jugular vein approach)

In recommending this procedure your doctor has balanced the benefits and risks of the procedure against the benefits and risks of not proceeding. Your doctor believes there is a net benefit to you going ahead. This is a very complicated assessment.

There are risks and complications with this procedure. They include but are not limited to the following.

Common risks and complications (more than 5%) include:

- Minor bleeding and bruising at the puncture site.
- Abnormal heartbeat lasting several seconds, which settles by itself.

Uncommon risks and complications (1- 5%) include:

• Unable to get the catheter into the neck vein. The procedure may be changed to a different approach eg an arm or a leg vein.

(Affix identification label here)

Family name:

URN:

No

No

No

No

Given name(s):

Address:

Date of birth:

 Abnormal heart rhythm that continues for a long time. This may need an electric shock to correct.

Sex: 🗆 M

E

1

• The carotid artery (in the neck) is accidentally punctured. This may require surgery to repair.

Rare risks and complications (less than 1%) include:

- Infection. This will need antibiotics.
- Allergic reaction to the local anaesthetic. This
 may require medication to treat.
- Blood clot in the neck vein. This may need medication to treat.
- Embolism. A blood clot may form and break off from the catheter. This is treated with blood thinning medication.
- Air in the lung cavity. A chest tube may need to be put in to the chest to drain the air.
- Damage to the vein in the neck causing bleeding. This may need surgery to repair.
- Air embolism. Oxygen may be given.
- A hole is accidentally made in the heart or the heart valve. This will need surgery to repair.
- Unable to position the balloon catheter into the lung vessels or around the heart. The procedure would be cancelled if this occurred. This is more common if there are congenital malformations of the heart.
- Damage to the lung blood vessel causing bleeding. This may need surgery to repair.
- A stroke. This may cause long term disability.
- Death as a result of this procedure is extremely rare.

D. Significant risks and procedure options

(Doctor to document in space provided. Continue in Medical Record if necessary.)

E. Risks of not having this procedure

(Doctor to document in space provided. Continue in Medical Record if necessary.)

F. Anaesthetic

This procedure may require an anaesthetic. (Doctor to document type of anaesthetic discussed)



Right Heart Catheter (Internal Jugular Vein Approach)

	(Affix iden	tification label	here)		
URN:					
Family name:					
Given name(s):					
Address:					
Date of birth:			Sex: M	F	
_	-	-	_		

Facility:

G. Patient consent

I acknowledge that the doctor has explained;

- my medical condition and the proposed procedure, including additional treatment if the doctor finds something unexpected. I understand the risks, including the risks that are specific to me.
- the anaesthetic required for this procedure. I understand the risks, including the risks that are specific to me.
- other relevant procedure/treatment options and their associated risks.
- my prognosis and the risks of not having the procedure.
- that no guarantee has been made that the procedure will improve my condition even though it has been carried out with due professional care.
- the procedure may include a blood transfusion.
- tissues and blood may be removed and could be used for diagnosis or management of my condition, stored and disposed of sensitively by the hospital.
- if immediate life-threatening events happen during the procedure, they will be treated based on my discussions with the doctor or my Acute Resuscitation Plan.
- a doctor other than the Consultant may conduct the procedure. I understand this could be a doctor undergoing further training.

I have been given the following Patient Information Sheet/s:

- Local Anaesthetic and Sedation for Your Procedure
 - Right Heart Catheter (Internal Jugular Vein Approach)
- I was able to ask questions and raise concerns with the doctor about my condition, the proposed procedure and its risks, and my treatment options. My questions and concerns have been discussed and answered to my satisfaction.
- I understand I have the right to change my mind at any time, including after I have signed this form but, preferably following a discussion with my doctor.
- I understand that image/s or video footage may be recorded as part of and during my procedure and that these image/s or video/s will assist the doctor to provide appropriate treatment.

On the basis of the above statements,

I request to have the procedure

Name of Patient:

Signature:

Date:

Consent	s who lack capacity to provide consent must be obtained from a substitute decision n the order below.
Does the (AHD)?	patient have an Advance Health Directive
□ Yes►	Location of the original or certified copy of the AHD:
□ No ►	Name of Substitute Decision Maker/s:
	Signature:
	Relationship to patient:
	Date: PH No:
	 Source of decision making authority (tick one): Tribunal-appointed Guardian Attorney/s for health matters under Enduring Power of Attorney or AHD Statutory Health Attorney If none of these, the Adult Guardian has provided consent.

H. Doctor/Delegate statement

I have explained to the patient all the above points under the Patient Consent section (G) and I am of the opinion that the patient/substitute decisionmaker has understood the information.

Name of

Doctor/delegate:

Designation:

Signature:

Date:

I. Interpreter's statement

I have given a sight translation in

(state the patient's language here) of the consent form and assisted in the provision of any verbal and written information given to the patient/parent or guardian/substitute decision-maker by the doctor. Name of

Interpreter:

Signature:

Date:

02/2011 - v5.00

1. What is a right heart catheter (internal jugular vein approach)?

This involves placing a fine tube (catheter), which will monitor pressure, into the chambers of your heart and the blood vessels of your lungs. By doing this, the doctors will learn more about the pressure in your heart and lungs and the pumping function of your heart.

You will have the following procedure:

A needle with a tube connected to it will be put in your arm. This is called an intravenous line or IV.

The vein in your neck is found with ultrasound.

You will be given an injection of local anaesthetic. A fine tube (catheter) is put into the vein in the neck. You may feel pressure on your neck while the tube is placed in the vein. It is passed down until it reaches the heart and then goes up into the blood vessels of the lungs. This is usually painless. The doctor uses x-ray imaging to see the catheter.

Pressures in the lungs and the heart are recorded. A sample of blood is taken to look at the oxygen level and other levels.

At the end of the procedure, the catheter is removed.

2. My anaesthetic

This procedure will require an anaesthetic.

See Local Anaesthetic and Sedation for Your Procedure information sheet for information about the anaesthetic and the risks involved. If you have any concerns, discuss these with your doctor.

If you have not been given an information sheet, please ask for one.

3. What are the risks of this specific procedure?

In recommending this procedure your doctor has balanced the benefits and risks of the procedure against the benefits and risks of not proceeding. Your doctor believes there is a net benefit to you going ahead. This is a very complicated assessment.

There are risks and complications with this procedure. They include but are not limited to the following.

Common risks and complications (more than 5%) include:

- Minor bleeding and bruising at the puncture site.
- Abnormal heartbeat lasting several seconds, which settles by itself.

Uncommon risks and complications (1- 5%) include:


- Unable to get the catheter into the neck vein. The procedure may be changed to a different approach eg an arm or a leg vein.
- Abnormal heart rhythm that continues for a long time. This may need an electric shock to correct.

• The carotid artery (in the neck) is accidentally punctured. This may require surgery to repair.

Rare risks and complications (less than 1%) include:

- Infection. This will need antibiotics.
- Allergic reaction to the local anaesthetic. This may require medication to treat.
- Blood clot in the neck vein. This may need medication to treat.
- Embolism. A blood clot may form and break off from the catheter. This is treated with blood thinning medication.
- Air in the lung cavity. A chest tube may need to be put in to the chest to drain the air.
- Damage to the vein in the neck causing bleeding. This may need surgery to repair.
- Air embolism. Oxygen may be given.
- A hole is accidentally made in the heart or the heart valve. This will need surgery to repair.
- Unable to position the balloon catheter into the lung vessels or around the heart. The procedure would be cancelled if this occurred. This is more common if there are congenital malformations of the heart.
- Damage to the lung blood vessel causing bleeding. This may need surgery to repair.

• A stroke. This may cause long term disability. Death as a result of this procedure is extremely rare.



Medical Illustration Copyright © 2008 Nucleus Medical Art, All rights reserved. www.nucleusinc.com

Notes to talk to my doctor about: